

1 f

1 1 f

B Liu¹, K-W Lee¹, M Anzo¹, B Zhang², Xi³, Yo⁴, L Siy⁵, M Pak⁴, S Lin⁶ and Chen¹

1

nyab a ftangiogenesis ; BP3is
anionalyupguedding hpa, a pent
inbr of angiogenesis (az-Gnzatz
20 On endhelaleb Koong , 20 and
BP3M is pdimanyepdin he
asahrendhelaleb of huan fir , 20
at fan , 19 and bine opsutea
Bow and Bach, 20) kih aggesa pib
inve n tin angiogenesis regulation, phasap
of a fe k k h a n i s i e r o p a b i n t e
hat BP3M is o r a b n a n y e p d
in h p i a - a s i a t e d n a n y a n g i o g e n e s i s
, 19 Lee , 19 and m e n d h e l a l e b
Sid , 20) n i y a r e n t p l a t o n
i c h i s B P 3 a s a n e y a n f a s i n h i l o r
i n d e h g a t e r e g u l a t o r o f a n g i o g e n e s i s i n h e a d a n d
n e k a m u s e l a i n o n d , 20
I t , w r o t h a t B P 3 h a s t h e E
i n d e p e n d e n t i n h i b i t o r o n a n g i o g e n e s i s I d
t o s e q u e a s p o f b o d e s t o s i e ,

egulation) in addition to primary actions, hereby
 regulates endothelial function and subsequent
 neovascularization in
 fiberinegative effect of BP on E
 regulates vascular function, using an established
 model of human endothelial dysfunction
 in a rat model, 2003
 has no effect on vascular function in his assay
 however, endothelial dysfunction is
 observed. Finally, endothelial
 dysfunction is only inhibited by BP
 treatment (Figure 4)
 To further investigate the effect of BP on E
 regulates vascular function and examine its effects
 on endothelial function and time
 in this study, human endothelial
 cells were treated with E,
 BP, Long E, and an analog with des

really is a high affinity to be
 time has been (E) and
 & (E) exposed to endothelial cells
 for 20 minutes post-treatment
 is significantly higher than that of
 the non-treated cells. These observations suggest that
 E is involved in the regulation of time

\$ %

;;< "0"9.0 7 54 2 4 9 ;;<
"2 " 4 5 0 .5+

99
9.4 4 4 4 0 0494
.44 ;;< 6.4.4" 0 0 53 0 0
4 0 5 53 \$# ' " 5 5 4 2
... .. + #52 " 4 2 .2 4 3 77 "
\$ ' 0 4 ;;< 4" .. 7 + " %
". 0 52 4 3 4" 4 64 7 6 9.0
9.. . = 4.4" 0 0 53 . 6 0 6 .

m-

BP3a injection of the opening into
the type of one-stage green fluorescent
protein (GFP) expression system
represents an integration of a GFP gene
into the promoter of the 2D and
utilized in the high-throughput screening of
antiangiogenic screening of
the vascular endothelial cells in green
fluorescence, the bodies can be easily realized
in levels in his angiogenic line. As shown in
Figure 7, injection of BP3a of the
resulted in a sharp increase in the trunk
and tail regions (Figure 7) in the presence of
filizaton) by the general angiogenesis
related factors and the remaining
vascular abnormalities, regional
sites are observed on the 10
the other observed effect on the
function of BP3 in Esqation, the
formation of BP3 with itself
in the high angiogenic BP3 in
and in the 6mle period in
expression of the remaining
inhibitory factor in the BP3
indicating that the effect of BP3 on vas-
culogenesis is independent of Esqation. The
apoptosis induced by the remaining
vascularized Zeb1 mutants as the non-
dominant mutant with loss in nothod
differentiation (Chahal, 2005), 10 as well as

0494 "#.2. 43" \$ 5 4 . I E '
9 4 4 1# 5" @ " 0 53 + B 49 0 4 .
6 4 4" 64 .377 " . " 4 .5
0 "4 .+

: 94 vascular formation matrigel assays
CE?D#0 . \$+ %'6 .2 " " 5 7 4 5
< 052 4 5" 5 \$:94 # .3" # E '
64 64 4 053 4 5 ;;< ?D;<
\$ 5 4 . I E ' " 9 " 0 4
54 3.5 4553 \$F .0 0. >. #
E ' . =6 2 4+ 74) 034 4 & 1#
4 53 7 2 . 7 5" 3 9.0 43
.4 04 . 6. 0 4" " 50 .02 7 0 6
\$:64. + %%'+

Materials
8 0 53 4 5 ;;< " ;;< 6 34 " 7 5
:B DF# 2+\$; ? E ' A 4" ".4 "
4 (%!##+ 0494 7 ;;< 7 34 ;;<:3 "
" 34 7704 0 2 7 4 7 5 3 .4
0 0 #<&0 .6.0 @5"3 .4 "3 44
" \$ "54 4 ' "2 44 5 3
5" .. ".0 3" 2 9 . \$ 44 " **'
.2049 37 4 =2 5 4.\$"4 4. 6'+ 3;<;<
" 4 334 4. 5 .4 5 ;;< 6
2 0 ." 7 5 E2.4 4 :0+\$ 0" B E '+
8 ;;< 6. 34 " 7 5 ; 2 2 \$ " " .4 '+
4 \$ +!5-5' 6.7 5 04 F 0 . 36
\$ "7 " E '+

Chicken embryo chorioallantoic membrane assay
" "7 4 " 6 4 .6 0 0 " "
0 0 53 . 64 404 .6 0 7 2 0"
% %%552 .40 4 ". .+ 74 !" . 7 0 3 4
K #! 4 &1# " . 7 5 4 0 . 0 4 %
" + m 7 ;;< " " 5. \$ 55'
6. 52 4" 4 # 7 " 9" 53 .+
5. "... 6 5" 3 ".004 7 %5 7 %+ K
54 0 . 64 + 74) " . 7 0 3 4
53 . "# .6 =5 " 7 4 7 5 4 7 6
3 " 9 .. 4 @ " 7 4 52 4" " .3 " ..04
50 .02 + F .. 7 5 4 0 . 0 4 6
." . 49 0 4 . \$# + **('+

Murine matrigel plug angiogenesis assay
..6.. " .4 64 73 " 9 .. .7 5
.30 4 . 4 .. 7 5 0 4

0". 7 4 B 4 5 " 5 7 < 3 " 2 4
.. 4 7 7@ 4 < 3 " + J Clin Endocrinol
Metab <., *) *%+
" < 3 > # 8 D .. 5 ?
 , \$**(+ ? .0 " 4 64 7 04 # " 0 .
 .. + Proc Natl Acad Sci USA :7,
(*) *+
> 4 527 > ; 9 00 D ; C >
 . , \$**(+ .5 . 64 7 04 :
 " 2 .44 0 0 . , 2 .2 04 9 .4 " + Science38,
!)!!+
6 < 4 =4 8#+\$%%'+:.
64 7 04 3 " 2 4 ".4 . .4 0
" 2 0 4 .+ J Clin Endocrinol Metab; ; ! ((!) * +
.. # # >B 8 3 .4 + \$%%'+
8 2 " .. 7 .. " . 9 J .0 4
3 @. .. + Arterioscler Thromb Vasc Biol3 , *) * +
F 7 .; A9 .4 C>+\$%%%'? .0 " 4 64
7 04 " 4 .7 5 64 7 04 3 4 4 4
=2 .. 7 . 64 7 04 3 " 2 4
" 9 .. " 4 0 .+ Endocrinology-6- ,
%!)%!&+
F 7 4 6 C F>+\$**!' + :.
64 7 04 3 " 2 4 .4 .. 3 4 0
" 4 0 . . 7 04 7 0 " .4+ Biochem
Biophys Res Commun 333 &() (+
F ; > 8 .. > 8 4 ; :
4 + \$%%'+ 4 2 = "
..4 C:< 2 3 4 + Cancer Biol Ther6,
%)%! +
D 9 .D 0# 4 ; # B # . + \$%%'+
: . 64 7 04 \$;<'. . A " 7
" . 4 " 9 25 4 7 4 3 @. 53 +
Int J Dev Biol 6<) %+
D 0 . ;< B 4 B 5 . + \$**'+
: . 64 7 04 3 " 2 4 =2 ..
. .4 04 " 4 9 4 0 2 4 4 9 .+
Endocrinology- , &) &+
D " BD F . F8 . 8 +
\$**!' + 8 4 7 " 4 ;;< .4 .. "
.0 4 3 ;;<: " ;< 3 4 +

Endothelial negative regulator of angiogenesis in
head and neck squamous carcinoma